

Connes-amenability and normal, virtual diagonals for measure algebras, II

Volker Runde*

Abstract

We prove that the following are equivalent for a locally compact group G : (i) G is amenable; (ii) $M(G)$ is Connes-amenable; (iii) $M(G)$ has a normal, virtual diagonal.

Keywords: locally compact group; group algebra; measure algebra; amenability; Connes-amenability; normal, virtual diagonal.

2000 *Mathematics Subject Classification:* 22D15, 43A10 (primary), 43A20, 43A60, 46E15, 46H25, 46M20, 47B47.

The result

A Banach algebra $\mathfrak{A}$ is called *dual* if it is a dual space such that multiplication in $\mathfrak{A}$ is separately w^* -continuous. A Banach bimodule over a dual Banach algebra is called *normal* if it is a dual space such that the module operations are separately w^* -continuous (see [6, 7, 8]).

Definition A dual Banach algebra $\mathfrak{A}$ is called *Connes-amenable* if, for every normal Banach $\mathfrak{A}$ -bimodule E , every w^* -continuous derivation $D: \mathfrak{A} \rightarrow E$ is inner.

The notion of Connes-amenability was introduced for von Neumann algebras in [5] (the name “Connes-amenability” seems to originate from [3]); it is equivalent to a number of important von Neumann algebraic properties such as injectivity, semidiscreteness, and being approximately finite-dimensional (see [7, Chapter 6] for a self-contained exposition and references to the original literature).

For arbitrary dual Banach algebras, Connes-amenability was first considered in [6], and in [8] it was shown that $M(G)$, the measure algebra of a locally compact group G , is Connes-amenable if and only if G is amenable.

For any dual Banach algebra $\mathfrak{A}$, let $\mathcal{L}_{w^*}^2(\mathfrak{A}, \mathbb{C})$ denote the separately w^* -continuous bilinear functionals on $\mathfrak{A}$. It is easy to see that the multiplication map $\Delta: \mathfrak{A} \otimes \mathfrak{A} \rightarrow \mathfrak{A}$ extends to $\mathcal{L}_{w^*}^2(\mathfrak{A}, \mathbb{C})^*$ as a continuous $\mathfrak{A}$ -bimodule homomorphism.

*Research supported by NSERC under grant no. 227043-00.

Definition Let $\mathfrak{A}$ be a dual Banach algebra. A *normal, virtual diagonal* for $\mathfrak{A}$ is an element $M \in \mathcal{L}_{w^*}^2(\mathfrak{A}, \mathbb{C})^*$ such that

$$a \cdot M = M \cdot a \quad \text{and} \quad a \Delta M = a \quad (a \in \mathfrak{A}).$$

If $\mathfrak{A}$ has a normal, virtual diagonal, then it is Connes-amenable ([1, 2]). The converse holds if $\mathfrak{A}$ is a von Neumann algebra ([2]). It is an open question — likely with a negative answer — if Connes-amenable and the existence of normal, virtual diagonals are equivalent for arbitrary dual Banach algebras.

In [6] and [8], we gave partial positive answers for $\mathfrak{A} = M(G)$:

- If G is compact, then $M(G)$ has a normal, virtual diagonal ([6, Proposition 5.2] and [8, Proposition 3.3]).
- If G is discrete, then $M(G) = \ell^1(G)$ is Connes-amenable if and only if it has a normal, virtual diagonal ([8, Corollary 5.4]).

In this note, we will prove (and thus extend [8, Theorem 5.3]):

Theorem *The following are equivalent for a locally compact group G :*

- (i) G is amenable.
- (ii) $M(G)$ is Connes-amenable.
- (iii) $M(G)$ has a normal, virtual diagonal.

The proof

For convenience, we quote the following well-known characterization of amenable locally compact groups (see [7, Lemma 7.1.1], for example):

Lemma *A locally compact group G is amenable if and only if there is a net $(f_\alpha)_\alpha$ of non-negative functions in the unit sphere of $L^1(G)$ such that*

$$\sup_{x \in K} \|\delta_x * f_\alpha - f_\alpha\| \rightarrow 0 \tag{*}$$

for each compact subset K of G .

Proof of the Theorem In view of [8, Theorem 5.3], it is sufficient to show that (i) implies (iii).

Let $(f_\alpha)_\alpha$ be a net as specified in the lemma. Define a net $(m_\alpha)_\alpha$ in $M(G \times G)$ by letting

$$\langle f, m_\alpha \rangle := \int_G f(x, x^{-1}) f_\alpha(x) dx \quad (f \in \mathcal{C}_0(G \times G)),$$

where dx denotes integration with respect to left Haar measure on G . Let $y \in G$, and note that, for $f \in \mathcal{C}_0(G \times G)$,

$$\begin{aligned} \langle f, (\delta_y \otimes \delta_e) * m_\alpha \rangle &= \int_G f(yx, x^{-1}) f_\alpha(x) dx \\ &= \int_G f(x, x^{-1}y) f_\alpha(y^{-1}x) dx \quad (\text{substitute } y^{-1}x \text{ for } x) \\ &= \int_G f(x, x^{-1}y) (\delta_y * f_\alpha)(x) dx \end{aligned}$$

and

$$\langle f, m_\alpha * (\delta_e \otimes \delta_y) \rangle = \int_G f(x, x^{-1}y) f_\alpha(x) dx.$$

It follows from (*) that

$$\sup_{y \in K} \|(\delta_y \otimes \delta_e) * m_\alpha - m_\alpha * (\delta_e \otimes \delta_y)\| \rightarrow 0 \quad (**)$$

for each compact subset K of G .

By [8, Proposition 3.1], we may identify the Banach $M(G)$ -bimodules $\mathcal{L}_{w*}^2(M(G), \mathbb{C})$ and

$$\mathcal{SC}_0(G \times G) := \{f \in \ell^\infty(G \times G) : f(\cdot, x), f(x, \cdot) \in \mathcal{C}_0(G) \text{ for each } x \in G\}.$$

Let $\mathcal{U}$ be an ultrafilter on the index set of $(m_\alpha)_\alpha$ that dominates the order filter. Define $M \in \mathcal{SC}_0(G \times G)^*$ by letting

$$\langle f, M \rangle := \lim_{\mathcal{U}} \int_{G \times G} f(x, y) dm_\alpha(x, y) \quad (f \in \mathcal{SC}_0(G \times G))$$

(since all functions in $\mathcal{SC}_0(G \times G)$ are measurable with respect to any Borel measure by [4], the integrals do exist). It is routinely seen that $\Delta M = \delta_e$.

Let $\mu \in M(G)$ and let $f \in \mathcal{SC}_0(G \times G)$. Then we have:

$$\begin{aligned} &|\langle f, \mu \cdot M - M \cdot \mu \rangle| \\ &= |\langle f \cdot \mu - \mu \cdot f, M \rangle| \\ &= \left| \lim_{\mathcal{U}} \int_{G \times G} \left(\int_G (f(zx, y) - f(x, yz)) d\mu(z) \right) dm_\alpha(x, y) \right| \\ &= \left| \lim_{\mathcal{U}} \int_G \left(\int_{G \times G} (f(zx, y) - f(x, yz)) dm_\alpha(x, y) \right) d\mu(z) \right| \quad (\text{by Fubini's theorem}) \\ &\leq \lim_{\mathcal{U}} \int_G \left| \int_{G \times G} (f(zx, y) - f(x, yz)) dm_\alpha(x, y) \right| d|\mu|(z) \\ &= \lim_{\mathcal{U}} \int_G \left| \int_{G \times G} f(x, y) d((\delta_z \otimes \delta_e) * m_\alpha - m_\alpha * (\delta_e \otimes \delta_z))(x, y) \right| d|\mu|(z) \\ &\leq \lim_{\mathcal{U}} \int_G \|f\| \|(\delta_z \otimes \delta_e) * m_\alpha - m_\alpha * (\delta_e \otimes \delta_z)\| d|\mu|(z) \\ &\rightarrow 0 \quad (\text{by } (**) \text{ and the inner regularity of } |\mu|). \end{aligned}$$

It follows that M is a normal, virtual diagonal for $M(G)$. $\square$

References

- [1] G. CORACH and J. E. GALÉ, Averaging with virtual diagonals and geometry of representations. In: E. ALBRECHT and M. MATHIEU (ed.s), *Banach Algebras '97*, pp. 87–100. Walter de Gruyter, 1998.
- [2] E. G. EFFROS, Amenability and virtual diagonals for von Neumann algebras. *J. Functional Analysis* **78** (1988), 137–153.
- [3] A. YA. HELEMSKIĬ, Homological essence of amenability in the sense of A. Connes: the injectivity of the predual bimodule (translated from the Russian). *Math. USSR-Sb* **68** (1991), 555–566.
- [4] B. E. JOHNSON, Separate continuity and measurability. *Proc. Amer. Math. Soc.* **20** (1969), 420–422.
- [5] B. E. JOHNSON, R. V. KADISON, and J. RINGROSE, Cohomology of operator algebras, III. *Bull. Soc. Math. France* **100** (1972), 73–79.
- [6] V. RUNDE, Amenability for dual Banach algebras. *Studia Math.* **148** (2001), 47–66.
- [7] V. RUNDE, *Lectures on Amenability*. Lecture Notes in Mathematics **1774**, Springer Verlag, 2002.
- [8] V. RUNDE, Connes-amenability and normal, virtual diagonals for measure algebras. *J. London Math. Soc.* **67** (2003), 643–656.

[February 1, 2008]

Address: Department of Mathematical and Statistical Sciences
University of Alberta
Edmonton, Alberta
Canada T6G 2G1

E-mail: `vrunde@ualberta.ca`

URL: `http://www.math.ualberta.ca/~runde/runde.html`